



#### Before your Implants

- We strongly recommend that you stop smoking 4 weeks prior to having dental Implants, Smoking can strongly impact the ability for the implant to fail or not.
- Hygiene is essential to the implant future and ensuring that a full hygiene routine is followed daily is essential, Brush minimum 2 times daily for two minutes and use mouth wash and Floss.
- If its possible for you to obtain a panoramic x-ray then please do so, this will allow us to give an accurate quote as we will know if Sinus lift or Bone Graft is required

#### **Guidelines for After Care of your new Implants**

- Take care to only have cold drinks and do not eat until the local anesthetic has worn off. Avoid hot drinks or hot food for the first day and do not 'swill' liquid over the area. Try not to disturb the area with your tongue or fingers. Do not undertake strenuous exercise for the first 48 hours (running / gym).
- You may have some swelling and/or bruising following your treatment - this usually reaches a peak 2 to 3 days later. This is quite normal, and both will subside naturally after a few days. Swelling can be reduced with ice packs (or a bag of frozen peas) wrapped in a towel. Hold on the cheek area for a maximum of 10 minutes at a time with 20 minutes break. After 24 hours gentle heat is more beneficial. Sleeping propped up slightly on 2-3 pillows may also help
- Pain should not be a big problem. Although you may be sore for a few days after any surgery in your mouth, this can be easily managed with simple analgesics (pain killers). You should take these regularly at the maximum stated dose for the first 2 days after your surgery. Take whatever painkillers you normally take for headaches, aches and sprains (ibuprofen and paracetamol make a good combination), and take your first dose before the local anesthetic has completely worn off.
- If after a few days you experience increasing pain and swelling, you must return to the clinic as soon as possible so that we can ensure you are not beginning to develop an infection.
- If you have a denture that covers the surgical area, please wear it as little as possible for the first week to protect the surgical site during its initial healing period. You should always leave the denture out at night.

- The stitches are dissolvable but often remain for around 2-3 weeks, if they are uncomfortable or annoying, you may contact the dentist to remove them before you travel home.
- Some minor bleeding after surgery in the mouth is normal. If this persists, apply pressure by biting firmly down over the area on a dampened gauze swab or clean handkerchief for 60 minutes whilst sitting upright. Do not keep checking or changing the gauze. You should contact us if bleeding persists for any reason after applying pressure in this way.
- If you have been given a course of antibiotics to take after your surgery, please ensure that you complete the course.
- On occasions the cover that is over the implants whilst its healing can fall out, this is nothing to worry about, if this does happen then extra cleaning must be taken around the area to avoid stuck food etc.
- very rarely implants can fall out, in most cases this is due to chewing or improper use, if this does happen, keep the implant safe for your next visit as losing it will incur charges to replace at approx. £400
- It is quite normal is the implant cover falls off during the healing process of the newly implanted Implant, if this happens extra care is to be taken around the area and extra cleaning and mouth wash should be used to ensure its kept clean.
- If anything changes prior to treatment but after we have assessed your images then you must inform us, failure to do so can result in extra work being required and further cost when at the clinic.

## **Cleaning**

- Successful oral surgery depends on keeping the mouth as clean as possible.
- Please start to use the mouthwash you have been given on the evening of your surgery and continue for 1 week. This is especially important. You should gently bathe the surgical site by holding approximately 15ml over the site (the equivalent of half a cap-full) for at least 1 minute, 3 times a day, for 7 days.
- You should also start cleaning your other teeth as normal with a toothbrush, starting on the evening of your surgery. Avoid the surgical site for the first few days, but then begin to carefully clean this area as well when tenderness permits.
- Hot salt mouthwashes (a cup of hot water with a teaspoon of salt) are very beneficial for healing in the first week but taste awful. Ensure that the mouthwash is not so hot that it scalds and then hold the hot mouthwash over the surgical site until it cools. Repeat as often as possible.
- Try to keep food away from the surgical area for as long as possible. Rinse following eating to keep the area clean.
- Smoking is NOT permitted until the wound has healed as this severely limits healing in the mouth.

- We want your recovery to be as smooth and pleasant as possible. It is vital to follow these instructions very carefully - if you have any concerns or questions regarding your progress, please do not hesitate to contact us where we will be happy to help

